

MEMORIAL UNIVERSITY OF NEWFOUNDLAND
DEPARTMENT OF MATHEMATICS AND STATISTICS

SECTION 1.4

Math 1000 Worksheet

FALL 2022

SOLUTIONS

1. (a) Direct substitution produces a $\frac{0}{0}$ indeterminate form, so we use the cancellation method:

$$\lim_{x \rightarrow 4} \frac{2x^2 - 7x - 4}{3x^2 - 14x + 8} = \lim_{x \rightarrow 4} \frac{(2x + 1)(x - 4)}{(3x - 2)(x - 4)} = \lim_{x \rightarrow 4} \frac{2x + 1}{3x - 2} = \frac{9}{10},$$

exactly as we deduced using the numerical approach in Question 2(a) of the Worksheet for Section 1.2.

- (b) Direct substitution produces a $\frac{0}{0}$ indeterminate form. Since this is a rational function, we use the cancellation method:

$$\lim_{x \rightarrow -1} \frac{3x^2 - 9x - 12}{x^3 + 7x^2 + 15x + 9} = \lim_{x \rightarrow -1} \frac{3(x + 1)(x - 4)}{(x + 3)^2(x + 1)} = \lim_{x \rightarrow -1} \frac{3(x - 4)}{(x + 3)^2} = \frac{-15}{4}.$$

This corroborates our guess in Question 2(c) of the Worksheet for Section 1.2.

- (c) Direct substitution produces a $\frac{0}{0}$ indeterminate form, so we use the cancellation method:

$$\lim_{t \rightarrow 2} \frac{t^2 - t - 6}{t^3 - 6t^2 + 12t - 8} = \lim_{t \rightarrow 2} \frac{(t + 3)(t - 2)}{(t - 2)^3} = \lim_{t \rightarrow 2} \frac{t + 3}{(t - 2)^2}.$$

Now direct substitution produces a $\frac{K}{0}$ form, so the limit does not exist. As $t \rightarrow 2$ from either the left or the right, $(t + 3)$ tends towards 5 (a positive number) while $(t - 2)^2$ becomes a small positive number (because the squares of non-zero real numbers are always positive). Hence

$$\lim_{t \rightarrow 2} \frac{t^2 - t - 6}{t^3 - 6t^2 + 12t - 8} = \infty.$$

- (d) In this case, direct substitution results in a $\frac{K}{0}$ form, so we know that the limit **does not exist**. As $x \rightarrow \frac{1}{2}$ from either side, $3x$ approaches $\frac{3}{2}$ (a positive number). From the left as $x \rightarrow \frac{1}{2}$, $(2x - 1)$ tends towards a small negative number, so

$$\lim_{x \rightarrow \frac{1}{2}^-} \frac{3x}{2x - 1} = -\infty.$$

From the right as $x \rightarrow \frac{1}{2}$, $(2x - 1)$ tends towards a small positive number, so

$$\lim_{x \rightarrow \frac{1}{2}^+} \frac{3x}{2x - 1} = \infty.$$

Because the one-sided limits do not agree, we cannot assign ∞ or $-\infty$ to the limit.

- (e) Direct substitution produces a $\frac{0}{0}$ indeterminate form. This is a quasirational function, so we use the rationalisation method:

$$\begin{aligned} \lim_{x \rightarrow -4} \frac{\sqrt{x+8}-2}{x+4} \cdot \frac{\sqrt{x+8}+2}{\sqrt{x+8}+2} &= \lim_{x \rightarrow -4} \frac{(x+8)-4}{(x+4)(\sqrt{x+8}+2)} \\ &= \lim_{x \rightarrow -4} \frac{x+4}{(x+4)(\sqrt{x+8}+2)} \\ &= \lim_{x \rightarrow -4} \frac{1}{\sqrt{x+8}+2} = \frac{1}{4}. \end{aligned}$$

- (f) Direct substitution produces a $\frac{0}{0}$ indeterminate form, so we use the rationalisation method:

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{h^2-h}{\sqrt{h+3}-\sqrt{3}} \cdot \frac{\sqrt{h+3}+\sqrt{3}}{\sqrt{h+3}+\sqrt{3}} &= \lim_{h \rightarrow 0} \frac{h(h-1)(\sqrt{h+3}+\sqrt{3})}{(h+3)-3} \\ &= \lim_{h \rightarrow 0} \frac{h(h-1)(\sqrt{h+3}+\sqrt{3})}{h} \\ &= \lim_{h \rightarrow 0} (h-1)(\sqrt{h+3}+\sqrt{3}) \\ &= -2\sqrt{3}. \end{aligned}$$

- (g) In this case, we simply need to use direct substitution:

$$\lim_{x \rightarrow 3} \frac{x-5}{\sqrt{2x+3}+1} = \frac{-2}{\sqrt{9}+1} = -\frac{1}{2}.$$

- (h) Direct substitution produces a $\frac{0}{0}$ indeterminate form. We can rid ourselves of the negative exponent in the numerator by multiplying both the numerator and the denominator by $(x+1)$:

$$\frac{12(x+1)^{-1}-2}{x^2-6x+5} = \frac{12-2(x+1)}{(x+1)(x^2-6x+5)} = \frac{-2(x-5)}{(x+1)(x-1)(x-5)}.$$

Now the limit can be rewritten as

$$\begin{aligned} \lim_{x \rightarrow 5} \frac{12(x+1)^{-1}-2}{x^2-6x+5} &= \lim_{x \rightarrow 5} \frac{-2(x-5)}{(x+1)(x-1)(x-5)} \\ &= \lim_{x \rightarrow 5} \frac{-2}{(x+1)(x-1)} = \frac{-2}{24} = -\frac{1}{12}. \end{aligned}$$

- (i) Direct substitution yields a $\frac{0}{0}$ indeterminate form. This function can be rewritten in the manner of a normal rational function, which means that we can then use the cancellation method:

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{\frac{1}{h^2+9} - \frac{1}{9}}{h} &= \lim_{h \rightarrow 0} \frac{\frac{9-(h^2+9)}{9(h^2+9)}}{h} = \lim_{h \rightarrow 0} \frac{1}{h} \cdot \frac{-h^2}{9(h^2+9)} \\ &= \lim_{h \rightarrow 0} \frac{-h}{9(h^2+9)} = \frac{0}{81} = 0. \end{aligned}$$

- (j) Direct substitution produces a $\frac{0}{0}$ indeterminate form. The presence of sine functions suggests that we should use the special trigonometric limit. First let's deal with the sine function in the numerator. We need a factor of $8x$ in the denominator, so we write

$$\begin{aligned}\lim_{x \rightarrow 0} \frac{\sin(8x)}{\sin(2x)} &= \lim_{x \rightarrow 0} \frac{\sin(8x)}{8x} \cdot \frac{8x}{\sin(2x)} = \lim_{x \rightarrow 0} \frac{\sin(8x)}{8x} \cdot \lim_{x \rightarrow 0} \frac{8x}{\sin(2x)} \\ &= 1 \cdot \lim_{x \rightarrow 0} \frac{8x}{\sin(2x)} = \lim_{x \rightarrow 0} \frac{8x}{\sin(2x)}.\end{aligned}$$

To deal with the remaining sine function, observe that we can factor 4 out of the numerator to obtain a factor of $2x$:

$$\lim_{x \rightarrow 0} \frac{\sin(8x)}{\sin(2x)} = 4 \lim_{x \rightarrow 0} \frac{2x}{\sin(2x)} = 4 \lim_{x \rightarrow 0} \frac{1}{\left(\frac{\sin(2x)}{2x}\right)} = 4 \cdot \frac{1}{1} = 4.$$

Alternatively, we could use the double-angle formula for sine to write

$$\sin(8x) = 2 \sin(4x) \cos(4x) = 4 \sin(2x) \cos(2x) \cos(4x),$$

so

$$\begin{aligned}\lim_{x \rightarrow 0} \frac{\sin(8x)}{\sin(2x)} &= \lim_{x \rightarrow 0} \frac{4 \sin(2x) \cos(2x) \cos(4x)}{\sin(2x)} \\ &= 4 \lim_{x \rightarrow 0} \cos(2x) \cos(4x) = 4(1)(1) = 4\end{aligned}$$

by direct substitution.

- (k) Direct substitution yields a $\frac{0}{0}$ indeterminate form, so we will use a special trigonometric limit. Observe that

$$\begin{aligned}\lim_{x \rightarrow 0} \frac{1 - \cos^2(x)}{x} &= \lim_{x \rightarrow 0} \frac{[1 - \cos(x)][1 + \cos(x)]}{x} \\ &= \lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x} \cdot \lim_{x \rightarrow 0} [1 + \cos(x)] \\ &= 0 \cdot 2 = 0.\end{aligned}$$

- (ℓ) Direct substitution produces a $\frac{0}{0}$ indeterminate form. We can use the special trigonometric limit:

$$\begin{aligned}\lim_{x \rightarrow 0} \frac{\sin(3x^2)}{x \sin(x)} &= \lim_{x \rightarrow 0} \frac{3x \sin(3x^2)}{3x^2 \sin(x)} = \lim_{x \rightarrow 0} 3 \cdot \frac{x}{\sin(x)} \cdot \frac{\sin(3x^2)}{3x^2} \\ &= 3 \lim_{x \rightarrow 0} \frac{x}{\sin(x)} \cdot \lim_{x \rightarrow 0} \frac{\sin(3x^2)}{3x^2}.\end{aligned}$$

Observe that as $x \rightarrow 0$, $3x^2 \rightarrow 0$ as well, so

$$\lim_{x \rightarrow 0} \frac{\sin(3x^2)}{x \sin(x)} = 3(1)(1) = 3.$$

(m) By direct substitution, we obtain

$$\lim_{x \rightarrow \pi} \frac{\tan\left(\frac{x}{4}\right)}{1 - \cos(x)} = \frac{\tan\left(\frac{\pi}{4}\right)}{1 - \cos(\pi)} = \frac{1}{1 - (-1)} = \frac{1}{2}.$$

(n) Direct substitution produces a $\frac{0}{0}$ indeterminate form. Because this problem involves secant functions, we need to rewrite it in terms of other trigonometric functions if we're to use a special trigonometric limit. In particular, recall that $\sec(\theta) = \frac{1}{\cos(\theta)}$ so we have

$$\begin{aligned} \lim_{\theta \rightarrow 0} \frac{1 - \sec(\theta)}{\theta \sec(\theta)} &= \lim_{\theta \rightarrow 0} \frac{1 - \frac{1}{\cos(\theta)}}{\frac{\theta}{\cos(\theta)}} = \lim_{\theta \rightarrow 0} \frac{\frac{\cos(\theta) - 1}{\cos(\theta)}}{\frac{\theta}{\cos(\theta)}} \\ &= \lim_{\theta \rightarrow 0} \frac{\cos(\theta) - 1}{\theta} = 0. \end{aligned}$$

(o) Observe that $|x - 2|$ changes its definition at $x = 2$:

$$|x - 2| = \begin{cases} -(x - 2) & \text{if } x < 2 \\ x - 2 & \text{if } x \geq 2. \end{cases}$$

Thus we need to examine the one-sided limits. From the left,

$$\lim_{x \rightarrow 2^-} \frac{|x - 2| - 2}{x} = \lim_{x \rightarrow 2^-} \frac{-(x - 2) - 2}{x} = \lim_{x \rightarrow 2^-} \frac{-x}{x} = \lim_{x \rightarrow 2^-} (-1) = -1.$$

From the right,

$$\lim_{x \rightarrow 2^+} \frac{|x - 2| - 2}{x} = \lim_{x \rightarrow 2^+} \frac{(x - 2) - 2}{x} = \lim_{x \rightarrow 2^+} \frac{x - 4}{x} = \frac{-2}{2} = -1.$$

Since the one-sided limits agree, we can conclude that

$$\lim_{x \rightarrow 2} \frac{|x - 2| - 2}{x} = -1.$$

(p) Although $x \rightarrow -2$, $|x - 2|$ does not change its definition at $x = -2$, so we can just substitute directly:

$$\lim_{x \rightarrow -2} \frac{|x - 2| - 2}{x} = \frac{|-4| - 2}{-2} = -1.$$

(q) We must check the one-sided limits, since $|x|$ changes its definition at $x = 0$. For $x < 0$, $|x| = -x$ so we can write

$$\lim_{x \rightarrow 0^-} \frac{x^2 - 4x}{7x - |x|} = \lim_{x \rightarrow 0^-} \frac{x^2 - 4x}{7x - (-x)} = \lim_{x \rightarrow 0^-} \frac{x^2 - 4x}{8x} = \lim_{x \rightarrow 0^-} \frac{x - 4}{8} = -\frac{1}{2}.$$

For $x > 0$, $|x| = x$ so we have

$$\lim_{x \rightarrow 0^+} \frac{x^2 - 4x}{7x - |x|} = \lim_{x \rightarrow 0^+} \frac{x^2 - 4x}{7x - x} = \lim_{x \rightarrow 0^+} \frac{x^2 - 4x}{6x} = \lim_{x \rightarrow 0^+} \frac{x - 4}{6} = -\frac{2}{3}.$$

Since the one-sided limits are not equal, we can conclude that the given limit does not exist.

2. (a) Since $f(x)$ changes its definition at $x = 1$, we must check the one-sided limits. From the left,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x^2 + 3x + 5) = 9.$$

From the right,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (7x - 2) = 5.$$

Since these are not equal, $\lim_{x \rightarrow 1} f(x)$ does not exist.

- (b) Again, $g(x)$ changes its definition at $x = 1$, so we must check the one-sided limits. From the left,

$$\lim_{x \rightarrow 1^-} g(x) = \lim_{x \rightarrow 1^-} (x^2 + 3x + 5) = 9.$$

From the right,

$$\lim_{x \rightarrow 1^+} g(x) = \lim_{x \rightarrow 1^+} (7x + 2) = 9.$$

Since the one-sided limits agree, we can conclude that $\lim_{x \rightarrow 1} g(x) = 9$ as well.

- (c) This time, $h(x)$ does not change its definition at $x = 1$, so we can simply write

$$\lim_{x \rightarrow 1} h(x) = \lim_{x \rightarrow 1} (7x - 2) = 5.$$

3. (a) We set the denominator equal to zero, so that

$$x^3 + 3x^2 - 9x + 5 = (x + 5)(x - 1)^2 = 0.$$

Hence the only possible vertical asymptotes are $x = -5$ and $x = 1$.

When $x = -5$, the numerator is $-54 \neq 0$, so we have a $\frac{K}{0}$ form. Hence $x = -5$ is a vertical asymptote. From the left as $x \rightarrow -5$, the denominator is a small negative number, so given that the numerator is also negative,

$$\lim_{x \rightarrow -5^-} \frac{5x - 4 - x^2}{x^3 + 3x^2 - 9x + 5} = \infty.$$

From the right as $x \rightarrow -5$, the denominator is a small positive number, so

$$\lim_{x \rightarrow -5^+} \frac{5x - 4 - x^2}{x^3 + 3x^2 - 9x + 5} = -\infty.$$

When $x = 1$, however, the numerator is zero, so we have to take the limit using the cancellation method:

$$\lim_{x \rightarrow 1} \frac{5x - 4 - x^2}{x^3 + 3x^2 - 9x + 5} = \lim_{x \rightarrow 1} \frac{-(x - 4)(x - 1)}{(x + 5)(x - 1)^2} = \lim_{x \rightarrow 1} \frac{4 - x}{(x + 5)(x - 1)}.$$

Now direct substitution produces a $\frac{K}{0}$ form (with $K = 3$) so $x = 1$ is a vertical asymptote after all. From the left as $x \rightarrow 1$, the denominator is a small negative number, so

$$\lim_{x \rightarrow 1^-} \frac{5x - 4 - x^2}{x^3 + 3x^2 - 9x + 5} = -\infty.$$

From the right as $x \rightarrow 1$, the denominator is a small positive number, so

$$\lim_{x \rightarrow 1^+} \frac{5x - 4 - x^2}{x^3 + 3x^2 - 9x + 5} = \infty.$$

(b) We set the denominator equal to zero, so that

$$5x - x^2 - 4 = -(x - 4)(x - 1) = 0.$$

Hence the only possible vertical asymptotes are $x = 4$ and $x = 1$.

When $x = 4$, the numerator is $81 \neq 0$, so we have a $\frac{K}{0}$ form. Hence $x = 4$ is a vertical asymptote. From the left as $x \rightarrow 4$, the denominator is a small positive number, so

$$\lim_{x \rightarrow 4^-} \frac{x^3 + 3x^2 - 9x + 5}{5x - 4 - x^2} = \infty.$$

From the right as $x \rightarrow 4$, the denominator is a small negative number, so

$$\lim_{x \rightarrow 4^+} \frac{x^3 + 3x^2 - 9x + 5}{5x - 4 - x^2} = -\infty.$$

When $x = 1$, however, the numerator is zero, so we take the limit using the cancellation method:

$$\begin{aligned} \lim_{x \rightarrow 1} \frac{x^3 + 3x^2 - 9x + 5}{5x - 4 - x^2} &= \lim_{x \rightarrow 1} \frac{(x + 5)(x - 1)^2}{-(x - 4)(x - 1)} \\ &= \lim_{x \rightarrow 1} \frac{(x + 5)(x - 1)}{4 - x} = 0. \end{aligned}$$

Because $\lim_{x \rightarrow 1} f(x)$ exists, $x = 1$ is not a vertical asymptote.

4. Using the inequality, we can write

$$-\cot(x) \leq \cot(x) \sin\left(\frac{1}{x}\right) \leq \cot(x)$$

if $\cot(x) > 0$ or

$$-\cot(x) \geq \cot(x) \sin\left(\frac{1}{x}\right) \geq \cot(x)$$

if $\cot(x) < 0$. Furthermore,

$$\lim_{x \rightarrow \frac{\pi}{2}} \cot(x) = \cot\left(\frac{\pi}{2}\right) = 0 \quad \text{and} \quad \lim_{x \rightarrow \frac{\pi}{2}} -\cot(x) = -\cot\left(\frac{\pi}{2}\right) = 0.$$

Thus, by the Squeeze Theorem, we conclude that $\lim_{x \rightarrow \frac{\pi}{2}} \cot(x) \sin\left(\frac{1}{x}\right) = 0$ as well.

(Note that if you're not comfortable evaluating a cotangent directly, you can always use the identity $\cot(x) = \frac{\cos(x)}{\sin(x)}$. Here, for instance, $\cot\left(\frac{\pi}{2}\right) = \frac{\cos\left(\frac{\pi}{2}\right)}{\sin\left(\frac{\pi}{2}\right)} = \frac{0}{1} = 0$.)

5. We know that

$$-1 \leq \cos\left(\frac{\pi}{2x}\right) \leq 1.$$

If we multiply all parts of the inequality by some $x > 0$, we get

$$-x \leq x \cos\left(\frac{\pi}{2x}\right) \leq x.$$

On the other hand, if $x < 0$, the same multiplication flips the direction of the inequalities, giving

$$x \leq x \cos\left(\frac{\pi}{2x}\right) \leq -x.$$

We can combine these two cases if we recall that $|x| = x$ for $x > 0$ and $|x| = -x$ for $x < 0$. Thus we have

$$-|x| \leq x \cos\left(\frac{\pi}{2x}\right) \leq |x|.$$

We know that $\lim_{x \rightarrow 0} |x| = 0$ and so

$$\lim_{x \rightarrow 0} -|x| = -\lim_{x \rightarrow 0} |x| = 0$$

as well. By the Squeeze Theorem, then, we also have

$$\lim_{x \rightarrow 0} x \cos\left(\frac{\pi}{2x}\right) = 0.$$