

MEMORIAL UNIVERSITY OF NEWFOUNDLAND  
DEPARTMENT OF MATHEMATICS AND STATISTICS

SECTION 4.2

Math 1000 Worksheet

FALL 2022

**SOLUTIONS**

1. (a) The function is a polynomial, so it is defined for all real numbers  $x$ . Differentiation gives

$$f'(x) = 12x^3 - 24x^2 + 12x = 12x(x - 1)^2,$$

$$f''(x) = 36x^2 - 48x + 12 = 12(x - 1)(3x - 1)$$

Setting  $f'(x) = 0$  gives  $x = 0$  and  $x = 1$ . Since  $f'(x)$  is always defined, these are our critical points. Setting  $f''(x) = 0$  gives  $x = 1$  and  $x = \frac{1}{3}$ . Since  $f''(x)$  is also always defined, these are the only hypercritical points. We use these values to construct the sign patterns depicted in Figure 1.



Figure 1: Sign patterns for Section 4.2, Question 1(a).

We can see that  $f(x)$  is increasing on  $0 < x < 1$  and  $x > 1$ , and decreasing on  $x < 0$ . There is a relative minimum at  $x = 0$  but  $x = 1$  is a saddle point.

The function is concave upward for  $x < \frac{1}{3}$  and  $x > 1$ , and concave downward for  $\frac{1}{3} < x < 1$ . Both  $x = \frac{1}{3}$  and  $x = 1$  are points of inflection.

- (b) Since  $1 + x^2 > 0$  for all  $x$ , and a logarithmic function is defined as long as its argument is positive, the domain of  $f(x)$  consists of all real numbers  $x$ . Differentiation gives

$$f'(x) = \frac{2x}{1 + x^2} \quad \text{and} \quad f''(x) = \frac{-2(x - 1)(x + 1)}{(x^2 + 1)^2}.$$

Setting  $f'(x) = 0$  gives  $x = 0$ . Since we cannot have  $(x^2 + 1)^2 = 0$ ,  $f'(x)$  is always defined, and so  $x = 0$  is the only critical point. Setting  $f''(x) = 0$  gives  $x = \pm 1$ . Again,  $f''(x)$  is always defined, so these are the only hypercritical points. We use these values to construct the sign pattern shown in Figure 2.



Figure 2: Sign patterns for Section 4.2, Question 1(b).

We can see that  $f(x)$  is increasing on  $x > 0$  and decreasing on  $x < 0$ . There is a relative minimum at  $x = 0$  but there are no local maxima.

Furthermore,  $f(x)$  is concave upward on  $-1 < x < 1$  and concave downward on  $x < -1$  and  $x > 1$ . Both  $x = 1$  and  $x = -1$  are inflection points.