MEMORIAL UNIVERSITY OF NEWFOUNDLAND

DEPARTMENT OF MATHEMATICS AND STATISTICS

Section 1.5

Math 1000 Worksheet

Fall 2025

SOLUTIONS

1. We must first expand the numerator and denominator:

$$f(x) = \frac{x(x+1)(1-x)}{(2x+3)^2} = \frac{-x^3 + x}{4x^2 + 12x + 9}.$$

The highest power of x in the denominator is x^2 , so

$$\lim_{x \to \infty} f(x) = \lim_{x \to \infty} \frac{-x^3 + x}{4x^2 + 12x + 9} \cdot \frac{\frac{1}{x^2}}{\frac{1}{x^2}} = \lim_{x \to \infty} \frac{-x + \frac{1}{x}}{4 + \frac{12}{x} + \frac{9}{x^2}}.$$

As $x \to \infty$, the numerator tends towards $-\infty$, while the denominator tends towards 4. Hence

$$\lim_{x \to \infty} f(x) = -\infty.$$

As $x \to -\infty$, the numerator tends towards ∞ , while the denominator again tends towards 4. Thus

$$\lim_{x \to -\infty} f(x) = \infty.$$

2. (a) Since f(x) is a rational function, we only need to take one of the limits at infinity. The highest power of x in the denominator is x^4 , so

$$\lim_{x \to \infty} f(x) = \lim_{x \to \infty} \frac{6x^3 - 6x^4}{2x^4 - x^2 + 1} \cdot \frac{\frac{1}{x^4}}{\frac{1}{x^4}} = \lim_{x \to \infty} \frac{\frac{6}{x} - 6}{2 - \frac{1}{x^2} + \frac{1}{x^4}} = \frac{0 - 6}{2 - 0 + 0} = -3.$$

Hence the only horizontal asymptote of f(x) is the line y = -3.

(b) Dividing by x^3 in the numerator and denominator gives

$$\lim_{x \to \infty} g(x) = \lim_{x \to \infty} \frac{6x^2 - 2x + 5}{7x^3 + x^{\frac{3}{2}}} \cdot \frac{\frac{1}{x^3}}{\frac{1}{x^3}} = \lim_{x \to \infty} \frac{\frac{6}{x} - \frac{2}{x^2} + \frac{5}{x^3}}{7 + \frac{1}{x^{\frac{3}{2}}}} = \frac{0 - 0 + 0}{7 + 0} = 0.$$

The only horizontal asymptote of g(x) is the line y = 0 (the x-axis).

(c) First we expand the numerator and denominator:

$$h(x) = \frac{(x+1)^3}{(4x^2+1)(2x-3)} = \frac{x^3+3x^2+3x+1}{8x^3-12x^2+2x-3}$$

Now we can see that the highest power of x in the denominator is x^3 , so dividing the numerator and denominator by x^3 gives

$$\lim_{x \to \infty} h(x) = \lim_{x \to \infty} \frac{1 + \frac{3}{x} + \frac{3}{x^2} + \frac{1}{x^3}}{8 - \frac{12}{x} + \frac{2}{x^2} - \frac{3}{x^3}} = \frac{1 + 0 + 0 + 0}{8 - 0 + 0 - 0} = \frac{1}{8}.$$

The only horizontal asymptote of h(x) is the line $y = \frac{1}{8}$.

3. (a) Observe that the highest power of x in the denominator is x, so for both limits we'll divide the numerator and denominator by x.

First we take the limit as $x \to \infty$, recalling that for x > 0, $x = \sqrt{x^2}$. Hence

$$\lim_{x \to \infty} \frac{x + \sqrt{4x^2 + 2}}{x - 7} \cdot \frac{\frac{1}{x}}{\frac{1}{x}} = \lim_{x \to \infty} \frac{1 + \frac{\sqrt{4x^2 + 2}}{\sqrt{x^2}}}{1 - \frac{7}{x}} = \lim_{x \to \infty} \frac{1 + \sqrt{4 + \frac{2}{x^2}}}{1 - \frac{7}{x}}$$
$$= \frac{1 + \sqrt{4 + 0}}{1 - 0} = 3.$$

Next we take the limit as $x \to -\infty$, recalling that for x < 0, $x = -\sqrt{x^2}$. So then

$$\lim_{x \to -\infty} \frac{x + \sqrt{4x^2 + 2}}{x - 7} \cdot \frac{\frac{1}{x}}{\frac{1}{x}} = \lim_{x \to -\infty} \frac{1 + \frac{\sqrt{4x^2 + 2}}{-\sqrt{x^2}}}{1 - \frac{7}{x}} = \lim_{x \to -\infty} \frac{1 - \sqrt{4 + \frac{2}{x^2}}}{1 - \frac{7}{x}}$$
$$= \frac{1 - \sqrt{4 + 0}}{1 - 0} = -1.$$

Thus the horizontal asymptotes of f(x) are y=3 and y=-1.

(b) The largest power of x outside the square root is x. The largest power of x inside the square root is x^2 , but we treat it as if its exponent were reduced by half, making it effectively x as well. Thus we proceed as if the largest power of x in the denominator is x, and this is what we will divide the numerator and the denominator by.

First let's take the limit as $x \to \infty$. Recalling that for x > 0, $x = \sqrt{x^2}$, we have

$$\lim_{x \to \infty} \frac{2x+1}{5x - \sqrt{9x^2 - 4}} \cdot \frac{\frac{1}{x}}{\frac{1}{x}} = \lim_{x \to \infty} \frac{2 + \frac{1}{x}}{5 - \frac{\sqrt{9x^2 - 4}}{x}} = \lim_{x \to \infty} \frac{2 + \frac{1}{x}}{5 - \frac{\sqrt{9x^2 - 4}}{\sqrt{x^2}}}$$
$$= \lim_{x \to \infty} \frac{2 + \frac{1}{x}}{5 - \sqrt{9 - \frac{4}{x^2}}} = \frac{2 + 0}{5 - \sqrt{9 - 0}} = 1.$$

For the limit as $x \to -\infty$, we follow the same procedure but note that since x < 0, $x = -\sqrt{x^2}$. Now we have

$$\lim_{x \to -\infty} \frac{2x+1}{5x-\sqrt{9x^2-4}} \cdot \frac{\frac{1}{x}}{\frac{1}{x}} = \lim_{x \to -\infty} \frac{2+\frac{1}{x}}{5-\frac{\sqrt{9x^2-4}}{x}} = \lim_{x \to -\infty} \frac{2+\frac{1}{x}}{5-\frac{\sqrt{9x^2-4}}{-\sqrt{x^2}}}$$
$$= \lim_{x \to -\infty} \frac{2+\frac{1}{x}}{5+\sqrt{9-\frac{4}{x^2}}} = \frac{2+0}{5+\sqrt{9-0}} = \frac{1}{4}.$$

Hence the two horizontal asymptotes are y = 1 and $y = \frac{1}{4}$.

(c) Again, the largest power of x outside the square root is x and the largest power inside the square root is effectively x as well.

First let's take the limit as $x \to \infty$. Recalling that for x > 0, $x = \sqrt{x^2}$, we have

$$\lim_{x \to \infty} \frac{2x+1}{5x - \sqrt{25x^2 - 4}} \cdot \frac{\frac{1}{x}}{\frac{1}{x}} = \lim_{x \to \infty} \frac{2 + \frac{1}{x}}{5 - \frac{\sqrt{25x^2 - 4}}{x}} = \lim_{x \to \infty} \frac{2 + \frac{1}{x}}{5 - \frac{\sqrt{25x^2 - 4}}{\sqrt{x^2}}}$$
$$= \lim_{x \to \infty} \frac{2 + \frac{1}{x}}{5 - \sqrt{25 - \frac{4}{x^2}}} = \frac{2 + 0}{5 - \sqrt{25 - 0}} = \frac{2}{0}.$$

Since this is a $\frac{K}{0}$ form, this limit does not exist.

For the limit as $x \to -\infty$, we follow the same procedure but note that since x < 0, $x = -\sqrt{x^2}$. Now we have

$$\lim_{x \to -\infty} \frac{2x+1}{5x - \sqrt{25x^2 - 4}} \cdot \frac{\frac{1}{x}}{\frac{1}{x}} = \lim_{x \to -\infty} \frac{2 + \frac{1}{x}}{5 - \frac{\sqrt{25x^2 - 4}}{x}} = \lim_{x \to -\infty} \frac{2 + \frac{1}{x}}{5 - \frac{\sqrt{25x^2 - 4}}{-\sqrt{x^2}}}$$
$$= \lim_{x \to -\infty} \frac{2 + \frac{1}{x}}{5 + \sqrt{25 - \frac{4}{x^2}}} = \frac{2 + 0}{5 + \sqrt{25 - 0}} = \frac{1}{5}.$$

Hence the only horizontal asymptotes is $y = \frac{1}{5}$.