MEMORIAL UNIVERSITY OF NEWFOUNDLAND DEPARTMENT OF MATHEMATICS AND STATISTICS

Section 1.3

Math 1000 Worksheet

Fall 2024

SOLUTIONS

1. (a) Using Basic Limit Property #2,

$$\lim_{x \to p} [f(x) - g(x)] = \lim_{x \to p} f(x) - \lim_{x \to p} g(x) = -5 - 4 = -9.$$

(b) Using Basic Limit Property #s 2 and 3,

$$\lim_{x \to p} [g(x) - 2f(x)] = \lim_{x \to p} g(x) - 2\lim_{x \to p} f(x) = 4 - 2(-5) = 14.$$

(c) Using Basic Limit Property #5,

$$\lim_{x \to p} \frac{f(x)}{g(x)} = \frac{\lim_{x \to p} f(x)}{\lim_{x \to p} g(x)} = \frac{-5}{4} = -\frac{5}{4}.$$

(d) Using Basic Limit Property #4 and the property for radicals of functions,

$$\lim_{x \to p} f(x) \sqrt{g(x)} = \lim_{x \to p} f(x) \cdot \sqrt{\lim_{x \to p} g(x)} = -5 \cdot \sqrt{4} = -10.$$

2. (a) By direct substitution, $\lim_{x\to 5}(x^2 - 9x - 3) = 5^2 - 9(5) + 3 = 25 - 45 + 3 = -17.$

- (b) By direct substitution, $\lim_{x \to -3} \frac{\sqrt{1-x}}{x} = \frac{\sqrt{4}}{-3} = -\frac{2}{3}$.
- (c) By direct substitution, $\lim_{h \to 0} \frac{\cos(h)}{2^h} = \frac{\cos(0)}{2^0} = 1.$
- (d) Observe that

$$|x-2| = \begin{cases} x-2 & \text{for } x \ge 2\\ -(x-2) & \text{for } x < 2. \end{cases}$$

Since this is a piecewise function which changes definition at x = 2, we must evaluate both the lefthand and righthand limits as $x \to 2$.

For the lefthand limit (where x < 2), we can write |x - 2| = -(x - 2), giving

$$\lim_{x \to 2^{-}} \frac{|x-2|}{|x-2|} = \lim_{x \to 2^{-}} \frac{-(x-2)}{|x-2|} = \lim_{x \to 2^{-}} -1 = -1.$$

For the righthand limit (where x > 2), we can write |x - 2| = x - 2, giving

$$\lim_{x \to 2^+} \frac{|x-2|}{x-2} = \lim_{x \to 2^+} \frac{x-2}{x-2} = \lim_{x \to 2^+} 1 = 1$$

Since the one-sided limits are not equal, we can conclude that $\lim_{x\to 2} \frac{|x-2|}{x-2}$ does not exist.

3. (a) Although f(x) is a piecewise function, its definition does not change at $x = \frac{\pi}{6}$, so we can use direct substitution. Since $f(x) = \cos(x)$ for all $x \le 0$,

$$\lim_{x \to -\frac{\pi}{6}} f(x) = \lim_{x \to -\frac{\pi}{6}} \cos(x) = \cos\left(-\frac{\pi}{6}\right) = \frac{\sqrt{3}}{2}.$$

(b) Since f(x) changes its definition at x = 0, we must consider the one-sided limits. Immediately to the left of x = 0, $f(x) = \cos(x)$ so

$$\lim_{x \to 0^{-}} f(x) = \lim_{x \to 0^{-}} \cos(x) = \cos(0) = 1.$$

Immediately to the right of x = 0, f(x) = 1 - 4x so

$$\lim_{x \to 0^+} f(x) = \lim_{x \to 0^+} (1 - 4x) = 1 - 4(0) = 1.$$

Since these are in agreement,

$$\lim_{x \to 0} f(x) = 1.$$

(c) Again, because f(x) changes its definition at x = 3, we must calculate the one-sided limits. Immediately to the left of x = 3, f(x) = 1 - 4x so

$$\lim_{x \to 3^{-}} f(x) = \lim_{x \to 3^{-}} (1 - 4x) = 1 - 4(3) = -11.$$

Immediately to the right of x = 3, $f(x) = \frac{9}{x}$ so

$$\lim_{x \to 3^+} f(x) = \lim_{x \to 3^+} \frac{9}{x} = \frac{9}{3} = 3$$

Since the one-sided limits are not equal, $\lim_{x\to 3} f(x)$ does not exist.

4. Since this is a piecewise function whose behaviour changes at x = -2, we must check the one-sided limits:

$$\lim_{x \to -2^{-}} f(x) = \lim_{x \to -2^{-}} k^2 x = -2k^2$$

and

$$\lim_{x \to -2^+} f(x) = \lim_{x \to -2^+} (4k - x) = 4k + 2.$$

If the limit exists, then these one-sided limits must be equal, so we set

$$-2k^{2} = 4k + 2$$
$$2k^{2} + 4k + 2 = 0$$
$$2(k+1)^{2} = 0,$$

and hence k = -1.

(Note that the second part of the definition of f(x) did not affect our workings because it applies only for x = -2, which has no effect on the limit as x approaches -2.)