MEMORIAL UNIVERSITY OF NEWFOUNDLAND DEPARTMENT OF MATHEMATICS AND STATISTICS

Section 4.2

Math 1000 Worksheet

Fall 2024

SOLUTIONS

1. (a) The function is a polynomial, so it is defined for all real numbers x. Differentiation gives

$$f'(x) = 12x^3 - 24x^2 + 12x = 12x(x-1)^2,$$

$$f''(x) = 36x^2 - 48x + 12 = 12(x-1)(3x-1)$$

Setting f'(x) = 0 gives x = 0 and x = 1. Since f'(x) is always defined, these are our critical points. Setting f''(x) = 0 gives x = 1 and $x = \frac{1}{3}$. Since f''(x) is also always defined, these are the only hypercritical points. We use these values to construct the sign patterns depicted in Figure 1.

Figure 1: Sign patterns for Section 4.2, Question 1(a).

We can see that f(x) is increasing on 0 < x < 1 and x > 1, and decreasing on x < 0. There is a relative minimum at x = 0 but x = 1 is a saddle point.

The function is concave upward for $x < \frac{1}{3}$ and x > 1, and concave downward for $\frac{1}{3} < x < 1$. Both $x = \frac{1}{3}$ and x = 1 are points of inflection.

(b) Since $1 + x^2 > 0$ for all x, and a logarithmic function is defined as long as its argument is positive, the domain of f(x) consists of all real numbers x. Differentiation gives

$$f'(x) = \frac{2x}{1+x^2}$$
 and $f''(x) = \frac{-2(x-1)(x+1)}{(x^2+1)^2}$.

Setting f'(x) = 0 gives x = 0. Since we cannot have $(x^2 + 1)^2 = 0$, f'(x) is always defined, and so x = 0 is the only critical point. Setting f''(x) = 0 gives $x = \pm 1$. Again, f''(x) is always defined, so these are the only hypercritical points. We use these values to construct the sign pattern shown in Figure 2.

Figure 2: Sign patterns for Section 4.2, Question 1(b).

We can see that f(x) is increasing on x > 0 and decreasing on x < 0. There is a relative minimum at x = 0 but there are no local maxima.

Furthermore, f(x) is concave upward on -1 < x < 1 and concave downward on x < -1 and x > 1. Both x = 1 and x = -1 are inflection points.