


## A Summary of Trigonometry

One way to think about the trigonometric functions is in terms of the unit circle (that is, the circle centred at the origin, of radius 1). If  $(x, y)$  is a point on the unit circle corresponding to an angle  $\theta$ , then

$$x = \cos(\theta) \quad \text{and} \quad y = \sin(\theta).$$

Note that we begin measuring  $\theta$  from the point  $(1, 0)$  (so this corresponds to the angle  $\theta = 0$ ). If  $\theta > 0$  then it is measured counterclockwise, and if  $\theta < 0$  then it is measured clockwise.



Furthermore, observe that both sine and cosine are periodic with period  $2\pi$ , so for any angle  $\theta$ , we have

$$\cos(\theta + 2\pi) = \cos(\theta) \quad \text{and} \quad \sin(\theta + 2\pi) = \sin(\theta).$$

Additionally, from the graphs of the cosine and sine functions, we can see that

$$\cos(-\theta) = \cos(\theta) \quad \text{and} \quad \sin(-\theta) = -\sin(\theta).$$

Finally, we can observe that

$$-1 \leq \sin(\theta) \leq 1 \quad \text{and} \quad -1 \leq \cos(\theta) \leq 1.$$

There are a number of useful relationships between the trigonometric functions (called identities). On the next page is a list of the most important.

## Basic Identities

- $\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)}$
- $\cot(\theta) = \frac{\cos(\theta)}{\sin(\theta)} = \frac{1}{\tan(\theta)}$
- $\sec(\theta) = \frac{1}{\cos(\theta)}$
- $\csc(\theta) = \frac{1}{\sin(\theta)}$
- $\tan(-\theta) = -\tan(\theta)$
- $\sin^2(\theta) + \cos^2(\theta) = 1$
- $\tan^2(\theta) + 1 = \sec^2(\theta)$
- $1 + \cot^2(\theta) = \csc^2(\theta)$

## Sum and Difference Identities

- $\cos(\alpha + \beta) = \cos(\alpha)\cos(\beta) - \sin(\alpha)\sin(\beta)$
- $\cos(\alpha - \beta) = \cos(\alpha)\cos(\beta) + \sin(\alpha)\sin(\beta)$
- $\sin(\alpha + \beta) = \sin(\alpha)\cos(\beta) + \cos(\alpha)\sin(\beta)$
- $\sin(\alpha - \beta) = \sin(\alpha)\cos(\beta) - \cos(\alpha)\sin(\beta)$

## Double and Half-Angle Identities

- $\cos(2\theta) = \cos^2(\theta) - \sin^2(\theta)$
- $\sin(2\theta) = 2\sin(\theta)\cos(\theta)$
- $\cos\left(\frac{\theta}{2}\right) = \pm\sqrt{\frac{1 + \cos(\theta)}{2}}$
- $\sin\left(\frac{\theta}{2}\right) = \pm\sqrt{\frac{1 - \cos(\theta)}{2}}$