

MEMORIAL UNIVERSITY OF NEWFOUNDLAND
DEPARTMENT OF MATHEMATICS AND STATISTICS

SECTION 2.7

Math 2000 Worksheet

FALL 2018

SOLUTIONS

1. (a) We may integrate with respect to x and y in either order, for instance:

$$\begin{aligned}\iint_D \frac{1}{\sqrt{16-x^2}} dA &= \int_{-2}^2 \int_0^7 \frac{1}{\sqrt{16-x^2}} dy dx = \int_{-2}^2 \left[\frac{y}{\sqrt{16-x^2}} \right]_0^7 dx \\ &= \int_{-2}^2 \frac{7}{\sqrt{16-x^2}} dx = \left[7 \arcsin \left(\frac{x}{4} \right) \right]_{-2}^2 = \frac{7\pi}{3}.\end{aligned}$$

- (b) We can write

$$D = \{(x, y) \mid 0 \leq x \leq 3, 0 \leq y \leq x\}.$$

Then the integral becomes

$$\begin{aligned}\iint_D \frac{1}{\sqrt{16-x^2}} dA &= \int_0^3 \int_0^x \frac{1}{\sqrt{16-x^2}} dy dx = \int_0^3 \left[\frac{y}{\sqrt{16-x^2}} \right]_0^x = \int_0^3 \frac{x}{\sqrt{16-x^2}} dx \\ &= \left[-\sqrt{16-x^2} \right]_0^3 = 4 - \sqrt{7}.\end{aligned}$$

2. (a)
$$\begin{aligned}\int_2^4 \int_1^{\sqrt{y}} x(y^2 - 5y) dx dy &= \int_2^4 \left[x^2(y^2 - 5y) \right]_1^{\sqrt{y}} dy = \int_2^4 \frac{1}{2}(y^2 - 5y)(y - 1) dy \\ &= \int_2^4 \left[\frac{1}{2}y^3 - 3y^2 + \frac{5}{2}y \right] dy = \left[\frac{1}{8}y^4 - y^3 + \frac{5}{4}y^2 \right]_2^4 = -11\end{aligned}$$

- (b) We have

$$\begin{aligned}\int_0^1 \int_0^{y^2} \frac{y}{x^2 + y^2} dx dy &= \int_0^1 \left[\frac{y}{y} \arctan \left(\frac{x}{y} \right) \right]_0^{y^2} dy = \int_0^1 \left[\arctan \left(\frac{x}{y} \right) \right]_0^{y^2} dy \\ &= \int_0^1 \arctan(y) dy = \left[y \arctan(y) - \frac{1}{2} \ln|1 + y^2| \right]_0^1 \\ &= \frac{\pi}{4} - \frac{1}{2} \ln(2),\end{aligned}$$

where the latter integral can be evaluated using integration by parts.

- (c) We have

$$\begin{aligned}\int_1^{\sqrt[4]{10}} \int_0^x y^2 \sqrt{x^4 - 1} dy dx &= \int_1^{\sqrt[4]{10}} \left[\frac{1}{3} y^3 \sqrt{x^4 - 1} \right]_0^x dx = \int_1^{\sqrt[4]{10}} \left[\frac{1}{3} x^3 \sqrt{x^4 - 1} \right] dx \\ &= \left[\frac{1}{18} (x^4 - 1)^{\frac{3}{2}} \right]_1^{\sqrt[4]{10}} = \frac{3}{2},\end{aligned}$$

where the second integral can be evaluated using u -substitution.

(d) We have

$$\begin{aligned} \int_{\frac{\pi}{2}}^0 \int_0^{\sin(x)} e^{\cos(x)} dy dx &= \int_{\frac{\pi}{2}}^0 \left[ye^{\cos(x)} \right]_0^{\sin(x)} dx = \int_{\frac{\pi}{2}}^0 \sin(x) e^{\cos(x)} dx \\ &= \left[-e^{\cos(x)} \right]_{\frac{\pi}{2}}^0 = 1 - e, \end{aligned}$$

where again we may use u -substitution to evaluate the final integral.

3. (a) The region of integration can be written as $0 \leq y \leq x$ and $0 \leq x \leq \sqrt{\pi}$. Hence the integral becomes

$$\begin{aligned} \int_0^{\sqrt{\pi}} \int_0^x \sin(x^2) dy dx &= \int_0^{\sqrt{\pi}} \left[y \sin(x^2) \right]_0^x dx = \int_0^{\sqrt{\pi}} x \sin(x^2) dx \\ &= \left[-\frac{1}{2} \cos(x^2) \right]_0^{\sqrt{\pi}} = 1. \end{aligned}$$

(b) The region of integration can be written as $0 \leq x \leq \sqrt{y}$ and $0 \leq y \leq 9$. Hence the integral becomes

$$\int_0^9 \int_0^{\sqrt{y}} x e^{y^2} dx dy = \int_0^9 \left[\frac{1}{2} x^2 e^{y^2} \right]_0^{\sqrt{y}} dy = \int_0^9 \frac{1}{2} y e^{y^2} dy = \left[\frac{1}{4} e^{y^2} \right]_0^9 = \frac{1}{4} e^{81} - \frac{1}{4}.$$

(c) The region of integration can be written as $0 \leq y \leq 2x$ and $0 \leq x \leq 2$. Hence the integral becomes

$$\begin{aligned} \int_0^2 \int_0^{2x} \frac{y}{x^3 + 1} dy dx &= \int_0^2 \left[\frac{y^2}{2(x^3 + 1)} \right]_0^{2x} dx = \int_0^2 \frac{2x^2}{x^3 + 1} dx \\ &= \left[\frac{2}{3} \ln |x^3 + 1| \right]_0^2 = \frac{4}{3} \ln(3). \end{aligned}$$

(d) The region of integration can be written as $0 \leq x \leq \sin(y)$ and $0 \leq y \leq \frac{\pi}{2}$. Hence the integral becomes

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \int_0^{\sin(y)} \sqrt{1 + \cos(y)} dx dy &= \int_0^{\frac{\pi}{2}} \left[x \sqrt{1 + \cos(y)} \right]_0^{\sin(y)} dy \\ &= \int_0^{\frac{\pi}{2}} \sin(y) \sqrt{1 + \cos(y)} dy \\ &= \left[-\frac{2}{3} [1 + \cos(y)]^{\frac{3}{2}} \right]_0^{\frac{\pi}{2}} = \frac{4\sqrt{2} - 2}{3}. \end{aligned}$$

4. (a) First we need to solve for the points of intersection of the two curves. We set

$$x^2 + 2x = 24 - x^2 \implies 2(x + 4)(x - 3) = 0$$

so $x = -4$ or $x = 3$. Then the region D is bounded by $-4 \leq x \leq 3$ and $x^2 + 2x \leq y \leq 24 - x^2$. (The order of the latter inequality may be easily checked by graphing, or by substitution of a value of x in the interval $(-4, 3)$, such as $x = 0$.) Then we have

$$\begin{aligned} A &= \iint_D dA = \int_{-4}^3 \int_{x^2+2x}^{24-x^2} dy dx \\ &= \int_{-4}^3 \left[y \right]_{x^2+2x}^{24-x^2} dx \\ &= \int_{-4}^3 [24 - 2x - 2x^2] dx \\ &= \left[24x - x^2 - \frac{2}{3}x^3 \right]_{-4}^3 \\ &= \frac{343}{3}. \end{aligned}$$

- (b) Again we begin by solving for the points of intersection. First we note that $y = 9 - 3x$ can also be written as $x = 3 - \frac{1}{3}y$. Then we have

$$\sqrt{9-y} = 3 - \frac{1}{3}y \implies 9 - y = 9 - 2y + \frac{1}{9}y^2 \implies y \left(\frac{1}{9}y - 1 \right) = 0$$

so $y = 0$ or $y = 9$. Thus the region R is defined by $0 \leq y \leq 9$ and $3 - \frac{1}{3}y \leq x \leq \sqrt{9-y}$. Then we have

$$\begin{aligned} A &= \iint_R dA = \int_0^9 \int_{\frac{1}{3}y}^{\sqrt{9-y}} dx dy \\ &= \int_0^9 \left[x \right]_{\frac{1}{3}y}^{\sqrt{9-y}} dy \\ &= \int_0^9 \left[\sqrt{9-y} - \frac{1}{3}y \right] dy \\ &= \left[-\frac{2}{3}(9-y)^{\frac{3}{2}} - \frac{1}{6}y^2 \right]_0^9 \\ &= \frac{9}{2}. \end{aligned}$$