

MEMORIAL UNIVERSITY OF NEWFOUNDLAND  
DEPARTMENT OF MATHEMATICS AND STATISTICS

---

SECTION 1.3

Math 2000 Worksheet

WINTER 2020

---

**SOLUTIONS**

1. (a) We have

$$s_1 = a_1 = -1$$

$$s_2 = a_1 + a_2 = -1 + \frac{3}{2} = \frac{1}{2}$$

$$s_3 = a_1 + a_2 + a_3 = -1 + \frac{3}{2} + \frac{5}{7} = \frac{17}{14}$$

$$s_4 = a_1 + a_2 + a_3 + a_4 = -1 + \frac{3}{2} + \frac{5}{7} + \frac{1}{2} = \frac{12}{7}$$

$$s_5 = a_1 + a_2 + a_3 + a_4 + a_5 = -1 + \frac{3}{2} + \frac{5}{7} + \frac{1}{2} + \frac{9}{23} = \frac{339}{161}$$

(b) Note that this series has  $i = 3$  as its first index, so

$$s_1 = a_3 = -\frac{1}{6}$$

$$s_2 = a_3 + a_4 = -\frac{1}{6} + \frac{1}{24} = -\frac{1}{8}$$

$$s_3 = a_3 + a_4 + a_5 = -\frac{1}{6} + \frac{1}{24} - \frac{1}{120} = -\frac{2}{15}$$

$$s_4 = a_3 + a_4 + a_5 + a_6 = -\frac{1}{6} + \frac{1}{24} - \frac{1}{120} + \frac{1}{720} = -\frac{19}{144}$$

$$s_5 = a_3 + a_4 + a_5 + a_6 + a_7 = -\frac{1}{6} + \frac{1}{24} - \frac{1}{120} + \frac{1}{720} - \frac{1}{5040} = -\frac{37}{280}$$

2. (a) We have

$$\lim_{i \rightarrow \infty} \left(-\frac{3}{7}\right)^{i+1} = \lim_{i \rightarrow \infty} -\frac{3}{7} \left(-\frac{3}{7}\right)^i = 0,$$

so the Divergence Test yields no conclusion.

(b) We have

$$\lim_{i \rightarrow \infty} \frac{i^3}{i(4i^2 - 5)} = \lim_{i \rightarrow \infty} \frac{i^3}{4i^3 - 5i} = \frac{1}{4},$$

so by the Divergence Test, this series is divergent.

(c) We have

$$\lim_{k \rightarrow \infty} \frac{2^k}{5^{k-1}} = 5 \lim_{k \rightarrow \infty} \left(\frac{2}{5}\right)^k = 0,$$

so we can draw no conclusion from the Divergence Test.

(d) We have

$$\lim_{i \rightarrow \infty} i \sin\left(\frac{1}{i}\right) = \lim_{i \rightarrow \infty} \frac{\sin\left(\frac{1}{i}\right)}{\frac{1}{i}} = 1,$$

so by the Divergence Test, this series is divergent.

3. First note that  $a_1 = s_1 = 3$ . For  $n > 1$ , we simply have

$$a_n = s_n - s_{n-1} = \left[5 - \frac{2}{n^2}\right] - \left[5 - \frac{2}{(n-1)^2}\right] = \frac{4n-2}{n^2(n-1)^2}.$$

Finally,

$$\sum_{i=1}^{\infty} a_i = \lim_{n \rightarrow \infty} s_n = \lim_{n \rightarrow \infty} \left[5 - \frac{2}{n^2}\right] = 5.$$

4. (a) Decomposing into partial fractions gives

$$\frac{1}{(i+4)(i+5)} = \frac{1}{i+4} - \frac{1}{i+5}.$$

Then we observe that this is a telescoping series with

$$s_n = \left(\frac{1}{4} - \frac{1}{5}\right) + \left(\frac{1}{5} - \frac{1}{6}\right) + \cdots + \left(\frac{1}{n+3} - \frac{1}{n+4}\right) + \left(\frac{1}{n+4} - \frac{1}{n+5}\right) = \frac{1}{4} - \frac{1}{n+5}.$$

Hence

$$\sum_{i=0}^{\infty} \frac{1}{(i+4)(i+5)} = \lim_{n \rightarrow \infty} \left(\frac{1}{4} - \frac{1}{n+5}\right) = \frac{1}{4}.$$

(b) Again, this is a telescoping series, with

$$s_n = [1^2 - 2^2] + [2^2 - 3^2] + \cdots + [(n-1)^2 - n^2] + [n^2 - (n+1)^2] = 1 - (n+1)^2.$$

Hence

$$\sum_{n=1}^{\infty} [n^2 - (n+1)^2] = \lim_{n \rightarrow \infty} s_n = \lim_{n \rightarrow \infty} [1 - (n+1)^2] = -\infty.$$

Therefore, the series diverges.

(c) Decomposing into partial fractions gives

$$\frac{2}{i(i-1)(i+1)} = \frac{1}{i-1} - \frac{2}{i} + \frac{1}{i+1}.$$

Then we observe that this is a telescoping series with

$$\begin{aligned} s_n &= \left(\frac{1}{1} - \frac{2}{2} + \frac{1}{3}\right) + \left(\frac{1}{2} - \frac{2}{3} + \frac{1}{4}\right) + \left(\frac{1}{3} - \frac{2}{4} + \frac{1}{5}\right) \\ &+ \cdots + \left(\frac{1}{n-2} - \frac{2}{n-1} + \frac{1}{n}\right) + \left(\frac{1}{n-1} - \frac{2}{n} + \frac{1}{n+1}\right). \end{aligned}$$

The pattern here is slightly more complicated: given any three consecutive parenthetical groupings, the term in the middle with a coefficient of  $-2$  is cancelled out by the combination of the last term in the preceding group and the first term in the following group. For instance, in the terms listed above, the  $-\frac{2}{3}$  in the second group is cancelled out by the  $+\frac{1}{3}$  in the first group and the  $+\frac{1}{3}$  in the third group. Studying this carefully, we see that the only terms which fail to cancel out completely in this manner are the  $\frac{1}{1} - \frac{2}{2}$  from the first group, the  $\frac{1}{2}$  from the second group, the  $\frac{1}{n}$  from the second-last group, and the  $-\frac{2}{n} + \frac{1}{n+1}$  from the last group. Therefore, we can write

$$s_n = \frac{1}{1} - \frac{2}{2} + \frac{1}{2} + \frac{1}{n} - \frac{2}{n} + \frac{1}{n+1} = \frac{1}{2} - \frac{1}{n} + \frac{1}{n+1},$$

and

$$\sum_{i=2}^{\infty} \frac{2}{i(i-1)(i+1)} = \lim_{n \rightarrow \infty} s_n = \lim_{n \rightarrow \infty} \left( \frac{1}{2} - \frac{1}{n} + \frac{1}{n+1} \right) = \frac{1}{2}.$$

5. (a) First we need to rewrite the given series as

$$\sum_{i=1}^{\infty} 4 \left( \frac{2}{3} \right)^i = \frac{8}{3} \sum_{i=1}^{\infty} \left( \frac{2}{3} \right)^{i-1}.$$

This is a geometric series with  $r = \frac{2}{3}$ , so

$$\sum_{i=1}^{\infty} 4 \left( \frac{2}{3} \right)^i = \frac{8}{3} \cdot \frac{1}{1 - \frac{2}{3}} = \frac{8}{3} \cdot 3 = 8.$$

(b) Rewriting gives

$$\sum_{i=0}^{\infty} \frac{3^i - 4^i}{3^i 4^i} = \sum_{i=0}^{\infty} \frac{3^i}{3^i 4^i} - \sum_{i=0}^{\infty} \frac{4^i}{3^i 4^i} = \sum_{i=0}^{\infty} \left( \frac{1}{4} \right)^i - \sum_{i=0}^{\infty} \left( \frac{1}{3} \right)^i.$$

Both of these series are geometric; the first has  $r = \frac{1}{4}$  and the second has  $r = \frac{1}{3}$ . So then

$$\sum_{i=0}^{\infty} \frac{3^i - 4^i}{3^i 4^i} = \frac{1}{1 - \frac{1}{4}} - \frac{1}{1 - \frac{1}{3}} = \frac{4}{3} - \frac{3}{2} = -\frac{1}{6}.$$

(c) Rewriting gives

$$\sum_{i=1}^{\infty} \left( \frac{2}{5} \right)^{3i} = \sum_{i=1}^{\infty} \left( \frac{8}{125} \right)^i = \frac{8}{125} \sum_{i=1}^{\infty} \left( \frac{8}{125} \right)^{i-1},$$

which is a geometric series with  $r = \frac{8}{125}$ . Then

$$\sum_{i=1}^{\infty} \left( \frac{2}{5} \right)^{3i} = \frac{8}{125} \cdot \frac{1}{1 - \frac{8}{125}} = \frac{8}{117}.$$

(d) Rewriting gives

$$\sum_{i=1}^{\infty} (-1)^{i-1} (0.2)^{i-1} = \sum_{i=1}^{\infty} (-0.2)^{i-1},$$

which is a geometric series with  $r = -0.2$ . Then

$$\sum_{i=1}^{\infty} (-1)^{i-1} (0.2)^{i-1} = \frac{1}{1 - (-0.2)} = \frac{1}{1.2} = \frac{5}{6}.$$

6. (a) Rewriting gives

$$\sum_{i=0}^{\infty} \frac{(x-6)^i}{4^i} = \sum_{i=0}^{\infty} \left( \frac{x-6}{4} \right)^i,$$

which is a geometric series with  $r = \frac{x-6}{4}$ . For this to converge, we require

$$-1 < \frac{x-6}{4} < 1 \implies -4 < x-6 < 4 \implies 2 < x < 10.$$

For these values of  $x$ ,

$$\sum_{i=0}^{\infty} \frac{(x-6)^i}{4^i} = \frac{1}{1 - \left(\frac{x-6}{4}\right)} = \frac{4}{10-x}.$$

(b) This is a geometric series with  $r = \sin(x)$ . It will converge when

$$-1 < \sin(x) < 1 \implies x \neq \frac{(2k+1)\pi}{2},$$

where  $k$  is an integer. For such values of  $x$ ,

$$\sum_{i=1}^{\infty} [\sin(x)]^{i-1} = \frac{1}{1 - \sin(x)}.$$

7. (a) First observe that

$$\begin{aligned} 0.042424242\dots &= 0.042 + 0.00042 + 0.0000042 + \dots \\ &= 0.042(1 + 0.01 + 0.0001 + \dots) \\ &= \sum_{i=1}^{\infty} 0.042(0.01)^{i-1} \\ &= \sum_{i=1}^{\infty} \frac{42}{1000} \left( \frac{1}{100} \right)^{i-1} \\ &= \frac{21}{500} \sum_{i=1}^{\infty} \left( \frac{1}{100} \right)^{i-1}, \end{aligned}$$

which is a geometric series with  $r = \frac{1}{100}$ . Then

$$0.042424242\dots = \frac{21}{500} \sum_{i=1}^{\infty} \left( \frac{1}{100} \right)^{i-1} = \frac{21}{500} \cdot \frac{1}{1 - \frac{1}{100}} = \frac{7}{165}.$$

(b) We can write

$$\begin{aligned}
 19.920920920 \dots &= 19 + 0.920 + 0.000920 + 0.00000920 + \dots \\
 &= 19 + 0.920(1 + 0.001 + 0.000001 + \dots) \\
 &= 19 + \sum_{i=1}^{\infty} 0.920(0.001)^{i-1} \\
 &= 19 + \frac{23}{25} \sum_{i=1}^{\infty} \left(\frac{1}{1000}\right)^{i-1}
 \end{aligned}$$

which is a geometric series with  $r = \frac{1}{1000}$ . Then

$$19.920920920 \dots = 19 + \frac{23}{25} \sum_{i=1}^{\infty} \left(\frac{1}{1000}\right)^{i-1} = 19 + \frac{23}{25} \cdot \frac{1}{1 - \frac{1}{1000}} = 19 + \frac{920}{999} = \frac{19901}{999}.$$

8. On the first bounce, the ball reaches a height of  $1(0.6) = 0.6$ . On the second bounce, the ball reaches a height of  $(0.6)(0.6) = (0.6)^2$ . On the third bounce, the ball reaches a height of  $(0.6^2)(0.6) = (0.6)^3$ . Continuing in this manner, it's clear that on the ball's  $i$ th bounce, it reaches a height of  $(0.6)^i$ . On each bounce, the ball travels up to its maximum height and then back down to the ground, for a distance travelled on the  $i$ th bounce of  $2(0.6)^i$ . Not forgetting the initial 1 metre drop, then, the ball's total distance travelled is

$$\begin{aligned}
 &1 + 2(0.6) + 2(0.6)^2 + 2(0.6)^3 + \dots \\
 &= 1 + 2(0.6)[1 + 0.6 + (0.6)^2 + \dots] \\
 &= 1 + 1.2 \sum_{i=1}^{\infty} (0.6)^{i-1} \\
 &= 1 + 1.2 \cdot \frac{1}{1 - 0.6} \\
 &= 1 + 3 \\
 &= 4.
 \end{aligned}$$

So the ball travels a total of 4 metres.