

MEMORIAL UNIVERSITY OF NEWFOUNDLAND

DEPARTMENT OF MATHEMATICS AND STATISTICS

SECTION 2.4

Math 2000 Worksheet

WINTER 2020

SOLUTIONS

1. Observe that

$$\begin{aligned} \frac{\partial z}{\partial x} &= \ln(x+2y) + \frac{x}{x+2y} & \frac{\partial z}{\partial y} &= \frac{2x}{x+2y} \\ \frac{dx}{dt} &= \cos(t) & \frac{dy}{dt} &= -\sin(t) \end{aligned}$$

so then

$$\begin{aligned} \frac{dz}{dt} &= \frac{\partial z}{\partial x} \frac{dx}{dt} + \frac{\partial z}{\partial y} \frac{dy}{dt} \\ &= \left[\ln(x+2y) + \frac{x}{x+2y} \right] \cos(t) + \left[\frac{2x}{x+2y} \right] [-\sin(t)] \\ &= \ln(x+2y) \cos(t) + \frac{x \cos(t)}{x+2y} - \frac{2x \sin(t)}{x+2y} \\ &= \ln(x+2y) \cos(t) + \frac{x \cos(t) - 2x \sin(t)}{x+2y}. \end{aligned}$$

2. We have

$$\begin{aligned} \frac{\partial z}{\partial u} &= \cos(u) \tan(v) & \frac{\partial z}{\partial v} &= \sin(u) \sec^2(v) & \frac{\partial u}{\partial x} &= 3 \\ \frac{\partial v}{\partial x} &= 1 & \frac{\partial u}{\partial y} &= 1 & \frac{\partial v}{\partial y} &= -1 \end{aligned}$$

and so

$$\begin{aligned} \frac{\partial z}{\partial x} &= \frac{\partial z}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial z}{\partial v} \frac{\partial v}{\partial x} \\ &= \cos(u) \tan(v)(3) + \sin(u) \sec^2(v)(1) \\ &= 3 \cos(u) \tan(v) + \sin(u) \sec^2(v) \\ \frac{\partial z}{\partial y} &= \frac{\partial z}{\partial u} \frac{\partial u}{\partial y} + \frac{\partial z}{\partial v} \frac{\partial v}{\partial y} \\ &= \cos(u) \tan(v)(1) + \sin(u) \sec^2(v)(-1) \\ &= \cos(u) \tan(v) - \sin(u) \sec^2(v). \end{aligned}$$

3. We have

$$\begin{aligned} \frac{\partial w}{\partial x} &= \frac{z}{\sqrt{1-y^2}} & \frac{\partial w}{\partial y} &= \frac{xyz}{(1-y^2)^{\frac{3}{2}}} & \frac{\partial w}{\partial z} &= \frac{x}{\sqrt{1-y^2}} \\ \frac{\partial x}{\partial r} &= 2r & \frac{\partial z}{\partial r} &= 4\theta e^{4r\theta} & \frac{\partial y}{\partial \theta} &= -\sin(\theta) \\ \frac{\partial z}{\partial \theta} &= 4r e^{4r\theta} & \frac{\partial y}{\partial r} &= \frac{\partial x}{\partial \theta} = 0 \end{aligned}$$

so that

$$\begin{aligned}
 \frac{\partial w}{\partial r} &= \frac{\partial w}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial r} + \frac{\partial w}{\partial z} \frac{\partial z}{\partial r} \\
 &= \frac{z}{\sqrt{1-y^2}}(2r) + \frac{xyz}{(1-y^2)^{\frac{3}{2}}}(0) + \frac{x}{\sqrt{1-y^2}}(4\theta e^{4r\theta}) \\
 &= \frac{2rz}{\sqrt{1-y^2}} + \frac{4x\theta e^{4r\theta}}{\sqrt{1-y^2}} \\
 &= \frac{2rz + 4x\theta e^{4r\theta}}{\sqrt{1-y^2}} \\
 \frac{\partial w}{\partial \theta} &= \frac{\partial w}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial \theta} + \frac{\partial w}{\partial z} \frac{\partial z}{\partial \theta} \\
 &= \frac{z}{\sqrt{1-y^2}}(0) + \frac{xyz}{(1-y^2)^{\frac{3}{2}}}[-\sin(\theta)] + \frac{x}{\sqrt{1-y^2}}(4re^{4r\theta}) \\
 &= -\frac{xyz \sin(\theta)}{(1-y^2)^{\frac{3}{2}}} + \frac{4xre^{4r\theta}}{\sqrt{1-y^2}} \\
 &= \frac{4x(1-y^2)re^{4r\theta} - xyz \sin(\theta)}{(1-y^2)^{\frac{3}{2}}}.
 \end{aligned}$$

4. We write

$$F(x, y) = \sin(x) + \cos(y) - \sin(x) \cos(y) - 7$$

so that

$$F_x(x, y) = \cos(x) - \cos(x) \cos(y) \quad \text{and} \quad F_y(x, y) = -\sin(y) + \sin(x) \sin(y),$$

and hence

$$\frac{dy}{dx} = -\frac{F_x(x, y)}{F_y(x, y)} = -\frac{\cos(x) - \cos(x) \cos(y)}{-\sin(y) + \sin(x) \sin(y)} = \frac{\cos(x) - \cos(x) \cos(y)}{\sin(y) - \sin(x) \sin(y)}.$$

5. We set

$$F(x, y, z) = x^2 - \sqrt{y} + z^2 - 2xyz$$

so that

$$F_x(x, y, z) = 2x - 2yz, \quad F_y(x, y, z) = -\frac{1}{2\sqrt{y}} - 2xz, \quad F_z(x, y, z) = 2z - 2xy.$$

Thus

$$\begin{aligned}
 \frac{\partial z}{\partial x} &= -\frac{F_x(x, y, z)}{F_z(x, y, z)} = -\frac{2x - 2yz}{2z - 2xy} = \frac{yz - x}{z - xy} \\
 \frac{\partial z}{\partial y} &= -\frac{F_y(x, y, z)}{F_z(x, y, z)} = -\frac{-\frac{1}{2\sqrt{y}} - 2xz}{2z - 2xy} = \frac{1 + 4xz\sqrt{y}}{4\sqrt{y}(z - xy)}.
 \end{aligned}$$

6. We set

$$F(x, y, z) = \sin^2(x) \cos(z) - \tan(z) - \sec(x) \csc(y)$$

so that

$$F_x(x, y, z) = 2 \sin(x) \cos(x) \cos(z) - \sec(x) \tan(x) \csc(y),$$

$$F_y(x, y, z) = \sec(x) \csc(y) \cot(y)$$

$$F_z(x, y, z) = -\sin^2(x) \sin(z) - \sec^2(z).$$

Hence

$$\frac{\partial z}{\partial x} = -\frac{F_x(x, y, z)}{F_z(x, y, z)} = \frac{2 \sin(x) \cos(x) \cos(z) - \sec(x) \tan(x) \csc(y)}{\sin^2(x) \sin(z) + \sec^2(z)}$$

$$\frac{\partial z}{\partial y} = -\frac{F_y(x, y, z)}{F_z(x, y, z)} = \frac{\sec(x) \csc(y) \cot(y)}{\sin^2(x) \sin(z) + \sec^2(z)}.$$